



Nucleoside Analogues




High Sensitivity: Fudr
 Partial Sensitivity: 5FU, MTX, Capecitabine, UFT, Raltitrexed, Pemetrexed
 No Sensitivity: Gemcitabine, Cytarabine, Fludarabine

Resistance Factors



Class I (cytotoxic Agents)

Activation of Caspase (especially 3 and 9) and cytochrom C re



Mr Hermann


Industrial Area of Florina, GR 53100 – Florina, Greece

Tel.: +30 23850 41950, 41951, 41960, 41961, Fax.: +30 23850 41931

Website: www.rgcc-group.com E-mail: papasotiriou.ioannis@rgcc-genlab.com


Class II (Immunostimulants/immunomodulators)

Immunostimulants / Immunomodulators release of Cytokins and increase of PBMC & NK



Class III (PK inhibitors)

Inhibitors of growth factors receptor inhibitors of EGFr,IGFr,VEGFr,PDGFr, FGFr, signal transduction pathways



CONCLUSION:It seems that this specific population of malignant cell have greater sensitivity in Amygdalin-(B17), in Artecina, in Ascorbic acid, in Bio D Mulsion NuMedica Micellized D3, in DCA (dichloroacetate), in Super Artemisinin, in Vitanox, in Virxcan, in Genistein, in Indol 3 Carbinol, in Naltrexone, in Paw-Paw, in Quercetin, in VasuStatin and less in Agaricus Blazei Murill, in Berberine, in BetaLoe, in Butyric Acid, in C-statin, in Graviola, in Honopure, in Lycopene, in Metformin, in OPC, in Okinawa Extracts, in Okinawa Nano, in Oleander Extract, in Poly-MVA, in ProsStay, in Salicinium, in Scullcap, in Synergy K, in Teavigo, in Ukrain, in Fucoidan, in Hucoidan, in Mistletoe, in RadoQOL, in Thymex, in Zyflamend, in Aromat8-PN, in Curcumin (turmeric), in Melatonin, in Nattokinase, in Resveratrol, in Salvestrol.

Sincerely,

Ioannis Papatotiriou MD., PhD
 Head of molecular medicine dpt. of
R.G.C.C.-RESEARCH GENETIC CANCER CENTRE LTD

DISCLAIMER

This study is known as an Ex-Vivo type study (testing the actual tumor stem cells of an individual outside their body). This test will tell us what natural substances will induce apoptosis via the cytochrome c (esp. caspase 3 & 9 pathways) after the tumor stem cells and a single product have been in contact, in a well plate for 48 hours. We have found this test to be very accurate over the past 10+ years and thousands of test. However, it cannot take into account the many combinations of natural substances or the physiological dynamics of each individual that are required for life. We are also aware that natural substances can have a wide variety of additional benefits that may assist healthy individuals, as well as those with cancer. Therefore, even if a product shows not to induce apoptosis, on this test, it most likely will have many other benefits especially when used in combination with other therapies your health care provider may use. This is when you must rely on the skill, knowledge and training of your health care provider and their years of clinical experience (successes and failures) with the many various combinations which they have found to work in a clinical setting. The body is a wonderful, magnificent, dynamic organism and very complex.